

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/82, 9/10, 15/54, C12Q 1/68, C12N 5/10, A01H 5/00		A1	(11) International Publication Number: WO 98/04725 (43) International Publication Date: 5 February 1998 (05.02.98)
(21) International Application Number: PCT/US97/13358 (22) International Filing Date: 30 July 1997 (30.07.97) (30) Priority Data: 60/023,314 31 July 1996 (31.07.96) US (71) Applicant: YALE UNIVERSITY [US/US]; Office of Cooperative Research, Suite 401, 246 Church Street, New Haven, CT 06510 (US). (72) Inventors: DELLAPORTA, Stephen, L.; 32 Leetes Island Road, Branford, CT 06405-3317 (US). CHEN, Jychian; 2 floor, #18-1, Alley 3, Lane 67, Section 2, Yen Jiou Yuan Road, Nankang 11529, Taipei (**). (74) Agents: ADLER, Reid, G. et al.; Morgan, Lewis & Bockius, LLP, 1800 M Street, N.W., Washington, DC 20036 (US).			(81) Designated States: AU, CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>

(54) Title: METHODS FOR ALTERING THE RATE OF PLANT DEVELOPMENT AND PLANTS OBTAINED THEREFROM



(57) Abstract

The present invention is based on the unexpected observation that DNA methylation, particularly at cytosine residues, regulates the rate of development of a plant. Based on this observation, the present invention provides methods of increasing or decreasing the rate of development of a plant by either increasing or decreasing the amount of methylated DNA found in the plant. The present invention further provides plants that have been altered such that their rate of maturation is either increased or decreased relative to the rate of maturation of a non-altered plant.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**METHODS FOR ALTERING THE RATE OF PLANT DEVELOPMENT AND
PLANTS OBTAINED THEREFROM**

This application claims the benefit of U.S. Provisional Application No. 60/023,314, filed July 31, 1996, the specification of which is hereby incorporated by reference in its entirety.

5 **STATEMENT OF RIGHTS TO INVENTIONS MADE UNDER
FEDERALLY SPONSORED RESEARCH**

This invention was made with governmental support under National Institutes of Health grant GM38148 and National Science Council grant 81-0203-B001-14. The Government has certain rights in the invention.

TECHNICAL FIELD

10 The present application is in the field of plant developmental biology and relates to methods for altering the rate at which a plant develops using molecular genetic techniques.

BACKGROUND ART

Plant genomes contain relatively large amounts of the modified nucleotide 5-methylcytosine (5^mC) (Y. Greenbaum, *et al.*, *Nature* 292: 850 (1981)). Despite
15 evidence implicating cytosine methylation in plant epigenetic phenomena, such as repeat-induced gene silencing (TIGS), cosuppression, and inactivation of transposable elements (F. F. Assaad, *et al.*, *Plant Mol. Biol.* 22: 1,057 (1993); C. Napoli, *et al.*,

Plant Cell 2: 279 (1990); J. Bender *et al.*, *Cell* 83: 725 (1995); P.S. Chomet, *et al.*, *Genetics* 138: 213 (1994); R.A. Martienssen, *et al.*, *Curr. Opin. Genet. Dev.* 5: 234 (1995); M.A. Matzke, *et al.*, *Plant Physiol.*, 107: 679-685 (1995)), the role of cytosine methylation in plant developmental processes is not clear.

5 In *Arabidopsis*, *ddm* mutants (decrease in DNA methylation) have been isolated with reduced levels of cytosine methylation in repetitive DNA sequences, although these mutations do not result in any detectable change in DNA methyltransferase enzymatic activity (A. Vongs, *et al.*, *Science*, 260: 1,926 (1993), T. Kakutani, *et al.*, *Nucleic Acids Res.* 23: 130 (1995)). After several generations of
10 self-pollination, *ddm* mutants exhibit a slight delay (1.7 days) in flowering, altered leaf shape, and an increase in cauline leaf number (T. Kakutani, *et al.* (1995)).

The exact mechanisms that mediate plant development are presently not well understood. Plants that have an increased rate of development would be highly useful in plant breeding programs. Specifically, numerous plants, such as tree species, have
15 extremely long generation times and therefore the number of crosses that can be generated within a given year or plant cycle is limited. In one extreme case, certain species of bamboo flower only once every one hundred years. Methods which could be used to decrease the maturation time would be highly beneficial in breeding programs involving many plants.

20 A reduction in the rate a plant matures can be used to increase the biomass production of a given plant. For numerous plants, increases in biomass yields would

increase the economic value of the commercial plant. For example, flax, tobacco, alfalfa, spinach, lettuce, *etc.*

It is therefore the focus of the present invention to provide methods for increasing or decreasing the time required for a plant to mature as well as plants
5 which are produced by these methods.

All references disclosed throughout this application are hereby incorporated by reference in their entirety.

DISCLOSURE OF THE INVENTION

The present invention is based on the unexpected observation that a decrease
10 of about 70% in the amount of methylated DNA present in a plant genome results in a plant that requires more time to mature while an increase in the amount of methylated DNA present in a plant genome results in a plant that requires less time to mature.

Based on these observation the present invention provides methods of altering a plant, plant cells, plant tissues or plant seeds, so as to obtain a plant that has an altered rate
15 of maturation. In one method, the rate of maturation is increased by altering the plant, plant cells, plant tissues or plant seeds, using molecular techniques, such that the plant has a sufficient increase in methylated DNA so as to yield a plant that matures faster than a non-altered plant. In another embodiment, the rate of maturation is decreased by altering a plant, plant cells, plant tissues or plant seeds, using molecular
20 techniques, such that the plant has a sufficient decrease in methylated DNA so as to yield a plant that matures slower than a non-altered plant.

The present invention further provides plants that have an altered rate of maturation that have been produced using the methods herein described.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. The predicted *MET1* gene product and antisense construct.

5 A diagrammatic representation of the predicted gene product of the *MET1* locus is provided. The *MET1* protein is a 1,534-amino acid protein with a high degree of homology to the mouse MTase, particularly in the catalytic and NH₂-terminal foci targeting domains (E. J. Finnegan, *et al.*, *Nucleic Acids Res.* 21: 2,383 (1993)). The *MET1* antisense construct is shown in the bottom of the figure. See Example 1.

10 Figure 2. Southern analysis of repetitive and single-copy DNA methylation patterns.

Total genomic DNA (3 µg/lane) from antisense lines, wild-type, and the *ddm1* mutant were digested with *Hpa II* (left panels) or *Msp I* (right panels), subjected to electrophoresis in 0.8% agarose, and transferred to Zeta-Probe membranes (Bio-Rad) and hybridized to probes as follows: (Upper panels) filters were probed with a
15 180-bp centromere repeat (J. M. Martinez-Zapater, *et al.*, *Mol. Gen. Genet.*, 204: 417 (1985)) and 5S rDNA (B. R. Campbell, *et al.*, *Gene* 112: 225 (1992)); (Lower panels) filters were probed with four single-copy gene probes--
PHOSPHORIBOSYLANTHRANILATE TRANSFERASE 1 (PAT1), PROLIFERA
(PRL), CONSTITUTIVELY PHOTOMORPHOGENIC 1 (COP1), and ERECTA

(ER). Digestion of wild-type genomic DNA with *Eco RI* and *Bst NI*, isochizomers with differential sensitivity to cytosine methylation in the motif C^{5m}C(A/T)GG, showed only minor differences in wild-type DNA when probed with the centromere repeat and 5S rDNA probes suggesting C(A/T)G methylation may not be prevalent-shown by differential *Hpa II*-*Msp I* digestion of wild-type DNA to contain C^{5m}CGG methylation M. J. Ronemus, *et al.*, data not shown. The panel on the right was hybridized to a control gene, *LEAFY* (LFY) (D. Weigel, *et al.*, *Cell* 59: 843 (1992)), shown not to be methylated in wild-type DNA.

Figure 3. Southern and phenotypic analyses of Tr246 outcross progeny.

10 Genomic DNAs (4 µg/lane) from outcross progeny of strong antisense line Tr246 to wild-type *Arabidopsis* (Columbia strain with no mutations) were digested with *Hpa II* (Fig. 3 upper panels) or *Eco RI* (Fig. 3 lower panels). Southern analyses were done as described above. Filters were probed with the 180-bp centromere repeat (Martinez-Zapater *et al.* (1985) (Fig. 3 upper panels) or the CaMV 35S promoter
15 fragment from pMON530 (Fig. 3 lower panels). Symbols indicate the presence (+) or absence (-) of the antisense transgene. Phenotypic data for each individual plant are shown below each lane. Plants were grown under continuous light at 21°C.

GENERAL DESCRIPTION
BEST MODE FOR CARRYING OUT THE INVENTION

General Description

The present invention is based on the unexpected observation that DNA

5 methylation, particularly at cytosine nucleotides, is involved in regulating the rate of
plant development and maturation, particularly the rate at which flower producing
organs and structures mature. Specifically, the present invention is based on the
observation that decreasing the amount of methylated DNA results in plants that
mature at a slower rate than plants containing normal amounts of methylated DNA

10 while increasing the amount of methylated DNA leads to plants that mature at a faster
rate than plants with normal amounts of methylated DNA.

Based on these observations, the present invention provides methods for
altering the rate of maturation of a plant which comprises the step of genetically
altering the plant, using molecular techniques, so the plant has an altered degree of

15 DNA methylation when compared to a non-altered plant sufficient to alter the rate at
which the plant develops.

As used herein, maturation refers to the process of plant differentiation leading
to the production of flowers and other reproductive tissues. The rate of maturation is
said to be altered when the rate at which the plant develops flowers is either increased

20 or decreased relative to a non-altered plant. The degree that the rate of maturation is

altered will vary from plant to plant as well as between plants that have been altered using different methods.

In a preferred embodiment, the method will produce plants with an increased rate of maturation of about 10% to about 25% faster, more preferably about 25% to
5 about 50% faster, than the normal rate of maturation. For example, in *Arabidopsis*, flowers typically develop after 26 days under long day conditions. In the Examples, *Arabidopsis* plants were obtained which developed flowers after 10 to 12 days.

The result of accelerating plant development is the ability to decrease the generational time in plant breeding programs. Plants that have been altered using the
10 methods herein described for decreasing maturation times can be used in any conventional breeding program, producing accelerated results.

In another preferred embodiment, the methods of the present invention will produce plants with a decreased rate of maturation that is about 10% to about 25% slower, more preferably about 25% to about 50% slower, most preferably about 50%
15 to about 100% or more slower than the normal rate of maturation. In the Examples, altered *Arabidopsis* plants were obtained that matured and produced flowers at 45-47 days, representing an almost 100% increase in the days required for maturation when compared to non-altered plants.

When used to slow the rate of maturation, the present methods produce plants
20 that have increased biomass. As used herein, biomass refers to the total plant weight,

particularly leafy material. The amount of leaf material is increased in plants altered to have a slower rate of maturation.

Another consequence of delaying flower production is to produce plants that have more secondary branches and axial nodes. As a result, the plant, when it
5 matures, produces a larger number of flower organs than does the non-altered plant.

Any plant that can be genetically altered using molecular techniques and any plants propagated containing the genetic alteration can be used in the present method. Methods known in the art for molecularly altering a plant are discussed in detail below. The preferred plants include both dicot and monocotyledonous plants. The
10 most preferred plants are plants with economic value as a food or biomass source, or plants with long maturation times. Such plants include, but are not limited to, leafy plants such as tobacco, spinach, lettuce, and seed bearing plants such as rice, corn, soy bean *etc.*

The methods of the present invention rely on altering plants using molecular
15 techniques. As used herein, molecular techniques exclude classical genetic techniques such as breeding/selection, identification of random mutagenesis and chemical mutagenesis techniques. Molecular techniques refers to procedures in which DNA is manipulated in a test tube during at least one stage of the process, such as the direct manipulation of DNA or the use of shuttle host such as bacterium. Such methods are
20 well known in the art and are described in, for example, Sambrook, *et al.*, *Molecular Cloning: a Laboratory Manual*, Cold Spring Harbor Press (1989). Some of the

techniques that are used to alter a plant are discussed in more detail below.

Molecular techniques are differentiated from classical genetics in which randomly occurring spontaneous mutants or classical mutagenic techniques are applied to a given plant type.

5 The requirement of the use of molecular techniques is to avoid the present invention reading on altered plants presently known in the art that have been generated through non-molecular techniques such as random mutation. Such plants may be known to have an altered rate of development but the underlying mechanism was not known prior to the present invention. Based on the present invention, it is
10 likely that such plants will be found to have alterations in DNA methylation.

 Although the present invention is based on the use of molecular techniques, a skilled artisan can now employ a new selection criteria when altering plants using non-molecular methods; namely selecting plants generated through methods such as chemical mutagenesis for increased or decreased DNA methylation. Specifically,
15 plants and plant cells can be subjected to chemical mutagenesis, physical mutagenesis and the like and the resulting plant selected based on alterations in amount of DNA methylation or in the activity of the DNA methyl transferase. Plants can then be further propagated from the isolated and identified variants as having altered amounts of DNA that then correlates to an altered rate of maturation.

Methods to Decrease the Rate of Maturation

As provided above, the methods of the present invention can be used to alter plants such that the rate of maturation is decreased, thus producing plants that require more time to mature. To obtain a plant that has a reduced rate of maturation when
5 compared to a non-altered plant, molecular techniques are used to reduce the amount of methylated DNA present in the altered plant.

As used herein, a plant is said to have a reduced or decreased amount of methylated DNA when the plant has less methylated DNA than the non-altered plant. Of the four nucleotides, cytosine has been seen to be methylated in plants. The
10 present methods are therefore accomplished by targeting the methylation of cytosine.

What is contemplated is a reduction sufficient and effective to alter the rate of maturation in a manner useful for the purposes outlined herein. Thus any alteration in the amount of DNA methylation, so long as it results in an altered rate of development, is contemplated by the present invention.

15 In the preferred embodiment, the method of the present invention will result in plants having about a 10% to about a 25% reduction, more preferable about a 25% to about a 50% reduction, most preferably about a 50% to about a 70% or greater reduction in the amount of methylated DNA present when compared to non-altered plants. In the Examples that follows, plants having a 75% reduction in methylated
20 cytosine were obtained.

A variety of targets, strategies and molecular techniques can be used by a skilled artisan to reduce the amount of methylated DNA present in a plant. For example, to obtain a reduction in the amount of methylated DNA, molecular techniques can be used to reduce the level of expression or inactivate the gene
5 encoding one or more DNA methyl transferases that are normally produced in the plant. The methyl transferase can be altered in a variety of ways by a skilled artisan so as to obtain a reduction in the amount of methylated DNA in the plant.

Any of a plant's DNA methyl transferases genes can be used as a target in the present method. The most preferred targets are the genes encoding a cytosine methyl
10 transferase. Examples of DNA methyl transferase genes known in the art include cytosine methyltransferase, such as the *MET1 Arabidopsis* gene herein used, the human and mouse analogs, and microbial methyl transferases such as bacterial GC methyl transferase (for example, see Finnegan *et al.*, (1995)).

In the examples that follow, an antisense DNA expression element was created
15 in which a fragment of a plant DNA methyl transferase gene was placed into an expression vector and inserted to a plant such that an mRNA was produced that is complimentary to mRNA encoding a plant DNA methyl transferase. A skilled artisan can readily follow this procedure with any DNA methyl transferase gene, or fragment thereof. Further, a skilled artisan can readily obtain a DNA methyl transferase gene
20 from any desired plant for use in the present methods. The preparation of antisense expression vectors is discussed in detail below.

An alternative strategy to decrease the amount of methylated DNA present in a plant is to create knockout mutants in the plant in which one or more of the plant's DNA methyl transferase genes are inactivated. This can be accomplished through the use of homologous recombination to insert stop codons or large DNA fragments
5 within the coding region of a plant DNA methyl transferase gene. Alternatively, transposon mediated mutagenesis can be used for the same purpose. The preparation of recombination vectors is discussed in detail below.

Plants that have been altered to contain a reduced amount of methylated DNA can be further altered so as to contain an expression unit that expresses a DNA methyl
10 transferase that is expressed or is functional under inducible conditions. Such plants are of additional value because of the ability to induce expression of a DNA methyl transferase gene, allowing one to activate or deactivate methyl transferase expression thus creating the ability to control alterations in the rate of maturation. The preparation of inducible expression vectors is discussed in detail below.

15 Methods to Increase the Rate of Maturation

As provided above, the methods of the present invention can be used to alter plants such that the rate of maturation is increased, thus producing plants that require less time to mature. To obtain a plant that has an increased rate of maturation when compared to a non-altered plant, molecular techniques are used to increase the amount
20 of methylated DNA present in the altered plant.

As used herein, a plant is said to have a reduced or decreased amount of methylated DNA when the plant has less methylated DNA than the non-altered plant. Of the four nucleotides, cytosine has been seen to be methylated in plants. The present methods are therefore accomplished by targeting the methylation of cytosine.

5 In the preferred embodiment, the method of the present invention will result in plants having about a 10% to about a 25% increase, more preferable about a 25% to about a 50% increase, most preferably about a 50% to about a 70% or greater increase in the amount of methylated DNA present when compared to non-altered plants. In the Examples that follows, plants having an increase in methylated cytosine were
10 obtained.

A variety of targets, strategies and molecular techniques can be used by a skilled artisan to increase the amount of methylated DNA present in the plant. For example, to obtain an increase in the amount of methylated DNA, molecular techniques can be used to increase the level of one or more DNA methyl transferases
15 in the plant cell. The methyl transferase can be altered in a variety of ways by a skilled artisan so as to obtain an increase in the amount of methylated DNA in the plant.

For example, DNA methyl transferase encoding expression units can be inserted into a plant genome using molecular techniques. Such methyl transferase
20 expression units can be controlled either by a constitutive promoter, providing continuous expression of the DNA methyl transferase gene, or using an inducible

promoter, allowing one to turn on or off the expression of DNA methyl transferase. By controlling the level at which the DNA methyl transferase is expressed, through the use of specific promoter sequences or by altering a plant to contain one or more additional copies of a DNA methyl transferase gene, the amount of methylated DNA
5 found in the plant can be dramatically increased.

Any DNA methyl transferase gene, or active fragment thereof, can be used to alter a plant such that the plant will contain an increased amount of methylated DNA, so long as the DNA methyl transferase gene can be modified so that it is expressed in a plant. In the examples that follow, the *MET1* DNA methyl transferase gene from
10 *Arabidopsis* was utilized. This gene, or its homologue from another plant, can readily be used by a skilled artisan in any plant system. The construction of expression units encoding a DNA methyl transferase is described in detail below.

Plants that have been altered to contain a increased amount of methylated DNA can be further altered so as to contain an expression unit that expresses a DNA
15 methyl transferase that is expressed or is functional under inducible conditions. Such plants are of additional value because of the ability to induce expression of a DNA methyl transferase gene, allowing one to activate or deactivate methyl transferase expression thus creating the ability to control alterations in the rate of maturation. The preparation of inducible expression vectors is discussed in detail below.

Plants with Altered Maturation Rates.

The present invention further provides plants that have been altered using molecular techniques so that they have an altered rate of maturation. As provided above, such plants will mature at a rate that is either slower than, or faster than, non-altered plants, depending on whether the amount of methylated DNA present is either increased or decreased relative to the amount present in a non-altered plant cell respectively. The plants of the present invention fall within two types. The first type are plants that have an increased rate of maturation while the second type are plants that have a reduced rate of maturation.

10 The preferred plants of the present invention that have an increased rate of maturation, will mature at a rate which is about 10% to about 25% faster, more preferably about 25% to about 50% faster, than the non-altered plant. The preferred plants of the present invention that have a slower rate of maturation will mature at a rate which is about 10% to about 25% slower, more preferably about 25% to about 15 50% slower, most preferably about 50% to about 100% slower than a non-altered plant.

The plants of the present invention include those that have been altered, using molecular techniques, to have an altered amount of methylated DNA sufficient to alter the rate of maturation of the plant. The plants of the present invention therefore 20 include any plant that can be altered using molecular technique so as to alter the amount of methylated DNA present in the plant. The preferred plants are plants for

which it is desirable to reduce generation times for breeding or seed production purposes or plants that are used for biomass production. Particularly useful plants include, but are not limited to, corn, rice, soy bean, wheat, spinach, lettuce, alfalfa, etc.

5 Expression Units to Express Exogenous DNA in a Plant

As provided above, several embodiments of the present invention employ expression units (or expression vectors or systems) to express an exogenously supplied gene, such as a DNA methyl transferase, or antisense molecule, in a plant. Methods for generating expression units/systems/vectors for use in plants are well known in the art and can readily be adapted for use in altering the amount of methylated DNA present in a plant cell. Typically, such units employ a protein or antisense coding region, such as the *MET1 Arabidopsis* gene, or a homologue thereof, and one or more expression control elements. The choice of the protein/antisense coding region, as well as the control elements, employed will be based on the effect desired (*i.e.*, reduction or increase in the amount of DNA methylation), the plant that is to be altered, the method chosen for altering the amount of methylated DNA and the transformation system used. A skilled artisan can readily use any appropriate plant/vector/expression system in the present methods following the outline provided herein.

10
15

The expression control elements used to regulate the expression of the protein or antisense coding region can either be the expression control element that is normally found associated with the coding sequence (homologous expression element) or can be a heterologous expression control element. A variety of

5 homologous and heterologous expression control elements are known in the art and can readily be used to make expression units for use in the present invention.

Transcription initiation regions, for example, can include any of the various opine initiation regions, such as octopine, mannopine, nopaline and the like that are found in the Ti plasmids of *Agrobacterium tumefaciens*. Alternatively, plant viral promoters

10 can also be used, such as the cauliflower mosaic virus 35S promoter to control gene expression in a plant. Lastly, plant promoters such as prolifera promoter, fruit-specific promoters, Ap3 promoter, heat shock promoters, seed-specific promoters, *etc.* can also be used. The most preferred promoters will be active in dividing tissue, particularly meristematic cells.

15 Either a constitutive promoter (such as the CaMV or Nos promoter illustrated above), an organ-specific promoter (such as the E8 promoter from tomato) or an inducible promoter is typically ligated to the protein or antisense encoding region using standard techniques known in the art. The expression unit may be further optimized by employing supplemental elements such as transcription terminators
20 and/or enhancer elements.

Thus, for expression in plants, the expression units will typically contain, in addition to the protein or antisense coding sequence, a plant promoter region, a transcription initiation site and a transcription termination sequence. Unique restriction enzyme sites at the 5' and 3' ends of the expression unit are typically included to allow for easy insertion into a preexisting vector. In the construction of heterologous promoter/structural gene or antisense combinations, the promoter is preferably positioned about the same distance from the heterologous transcription start site as it is from the transcription start site in its natural setting. As is known in the art, however, some variation in this distance can be accommodated without loss of promoter function.

In addition to a promoter sequence, the expression cassette can also contain a transcription termination region downstream of the structural gene to provide for efficient termination. The termination region may be obtained from the same gene as the promoter sequence or may be obtained from different genes. If the mRNA encoded by the structural gene is to be efficiently processed, DNA sequences which direct polyadenylation of the RNA are also commonly added to the vector construct. Polyadenylation sequences include, but are not limited to the *Agrobacterium* octopine synthase signal (Gielen *et al.*, *EMBO J* 3: 835-846 (1984)) or the nopaline synthase signal (Depicker *et al.*, *Mol. and Appl. Genet.* 1: 561-573 (1982)).

The resulting expression unit is ligated into or otherwise constructed to be included in a vector which is appropriate for higher plant transformation. The vector

will also typically contain a selectable marker gene by which transformed plant cells can be identified in culture. Usually, the marker gene will encode antibiotic resistance. These markers include resistance to G418, hygromycin, bleomycin, kanamycin, and gentamicin. After transforming the plant cells, those cells having the

5 vector will be identified by their ability to grow on a medium containing the particular antibiotic. Replication sequences, of bacterial or viral origin, are generally also included to allow the vector to be cloned in a bacterial or phage host, preferably a broad host range prokaryotic origin of replication is included. A selectable marker for bacteria should also be included to allow selection of bacterial cells bearing the

10 desired construct. Suitable prokaryotic selectable markers also include resistance to antibiotics such as kanamycin or tetracycline.

Other DNA sequences encoding additional functions may also be present in the vector, as is known in the art. For instance, in the case of *Agrobacterium* transformations, T-DNA sequences will also be included for subsequent transfer to

15 plant chromosomes.

Methods to produce antisense encoding vectors

As discussed above, plants having an altered amount of methylated DNA can be produced by using antisense sequences for interrupting the expression of one or more DNA methyl transferases in a plant cell. As evidenced by the behavior of

20 antisense mutants described in the Examples, reduction in DNA methylation results in

plants with reduced maturation rates while increases in DNA methylation results in plants with increased maturation rates. Accordingly, antisense sequences of suitable length can be transfected into plant cells, using the methods described herein, to obtain plants that take longer or shorter to mature than the non- altered plant.

- 5 Methods for inhibiting expression in plants using antisense constructs, including generation of antisense sequences *in situ* are described, for example, in U.S. Patents 5,107,065 and 5,254,800.

- Other methods that can be used to inhibit expression of an endogenous gene in a plant may also be used in the present methods. For example, formation of a triple
10 helix at an essential region of a duplex gene serves this purpose. The triplex code, permitting design of the proper single stranded participant is also known in the art. (See H. E. Moser, *et al.*, *Science* 238: 645-650 (1987) and M. Cooney, *et al.*, *Science* 241: 456-459 (1988)). Regions in the control sequences containing stretches of purine bases are particularly attractive targets. Triple helix formation along with
15 photocrosslinking is described, *e.g.*, in D. Praseuth, *et al.*, *Proc. Nat'l Acad. Sci. USA* 85: 1,349-1,353 (1988).

Inactivation of Endogenous Methyl transferases or transferases

- Another approach to inactivate one or more endogenous DNA plant genes that encode a DNA methyl transferase employs homologous recombination to disrupt the
20 gene. The techniques for recombinational inactivation are known in the art and can

readily be adapted to the present invention, for example see D. K. Asch, *et al.*, *Mol. Gen. Genet.* **221**: 37-43 (1990); K. K. Asch, *et al.*, *Genetics* **130**: 737-748 (1992).

Transformation of Plant Cells

When an appropriate vector is obtained, for example as described above,
5 transgenic plants are prepared which contain the desired expression unit or into which a recombination inactivation vector has been introduced. In one method of transformation, the vector is microinjected directly into plant cells by use of micropipettes to mechanically transfer the recombinant DNA into the plant cell (Crossway, *Mol. Gen. Genetics* **202**: 179-185 (1985)). In another method, the genetic
10 material is transferred into the plant cell using polyethylene glycol (Krens, *et al.*, *Nature* **296**: 72-74 (1982)), or high velocity ballistic penetration by small particles with the nucleic acid either within the matrix of small beads or particles, or on the surface, is used (Klein, *et al.*, *Nature* **327**: 70-73 (1987)). In still another method protoplasts are fused with other entities which contain the DNA whose introduction is
15 desired. These entities are minicells, cells, lysosomes or other fusible lipid-surfaced bodies (Fraley, *et al.*, *Proc. Nat'l Acad. Sci. USA* **79**: 1,859-1,863 (1982)).

DNA may also be introduced into the plant cells by electroporation (From *et al.*, *Proc. Nat'l Acad. Sci. USA* **82**: 5,824 (1985)). In this technique, plant protoplasts are electroporated in the presence of plasmids containing the expression cassette.

Electrical impulses of high field strength reversibly permeabilize biomembranes allowing the introduction of the plasmids. Electroporated plant protoplasts reform the cell wall, divide, and regenerate.

For transformation mediated by bacterial infection, a plant cell is infected with
5 *Agrobacterium tumefaciens* or *A. rhizogenes* previously transformed with the DNA to be introduced. *Agrobacterium* is a representative genus of the gram-negative family *Rhizobiaceae*. Its species are responsible for crown gall (*A. tumefaciens*) and hairy root disease (*A. rhizogenes*). The plant cells in crown gall tumors and hairy roots are induced to produce amino acid derivatives known as opines, which are catabolized
10 only by the bacteria. The bacterial genes responsible for expression of opines are a convenient source of control elements for chimeric expression cassettes. In addition, assaying for the presence of opines can be used to identify transformed tissue.

Heterologous genetic sequences can be introduced into appropriate plant cells, by means of the Ti plasmid of *A. tumefaciens* or the Ri plasmid of *A. rhizogenes*. The
15 Ti or Ri plasmid is transmitted to plant cells on infection by *Agrobacterium* and is stably integrated into the plant genome (J. Schell, *Science* 237: 1,176-1,183 (1987)). Ti and Ri plasmids contain two regions essential for the production of transformed cells. One of these, named transferred DNA (T-DNA), is transferred to plant nuclei and induces tumor or root formation. The other, termed the virulence (vir) region, is
20 essential for the transfer of the T-DNA but is not itself transferred. The T-DNA will be transferred into a plant cell even if the vir region is on a different plasmid

(Hoekema, *et al.*, *Nature* 303: 179-189 (1983)). The transferred DNA region can be increased in size by the insertion of heterologous DNA without its ability to be transferred being affected. Thus a modified Ti or Ri plasmid, in which the disease-causing genes have been deleted, can be used as a vector for the transfer of the

5 gene constructs of this invention into an appropriate plant cell.

Construction of recombinant Ti and Ri plasmids in general follows methods typically used with the more common bacterial vectors, such as pBR322. Additional use can be made of accessory genetic elements sometimes found with the native plasmids and sometimes constructed from foreign sequences. These may include but

10 are not limited to "shuttle vectors," (Ruvkun and Ausubel, *Nature* 298: 85-88 (1981)), promoters (Lawton *et al.*, *Plant Mol. Biol.* 9: 315-324 (1987)) and structural genes for antibiotic resistance as a selection factor (Fraley *et al.*, *Proc. Nat'l Acad. Sci.* 80: 4,803-4,807 (1983)).

There are two classes of recombinant Ti and Ri plasmid vector systems now in

15 use. In one class, called "co-integrate," the shuttle vector containing the gene of interest is inserted by genetic recombination into a non-oncogenic Ti plasmid that contains both the *cis*-acting and *trans*-acting elements required for plant transformation as, for example, in the pMLJ1 shuttle vector of DeBlock *et al.*, *EMBO J* 3: 1,681-1,689 (1984) and the non-oncogenic Ti plasmid pGV3850, described by

20 Zambryski *et al.*, *EMBO J* 2: 2,143-2,150 (1983). In the second class or "binary" system, the gene of interest is inserted into a shuttle vector containing the *cis*-acting

elements required for plant transformation. The other necessary functions are provided in *trans* by the non-oncogenic Ti plasmid, as exemplified by the pBIN19 shuttle vector described by Bevan, *Nucleic Acids Res.* 12: 8,711-8,721 (1984) and the non-oncogenic Ti plasmid pAL4404 described by Hoekma, *et al.*, (1983). Some of
5 these vectors are commercially available.

There are two common ways to transform plant cells with *Agrobacterium*: co-cultivation of *Agrobacterium* with cultured isolated protoplasts, or transformation of intact cells or tissues with *Agrobacterium*. The first requires an established culture system that allows for culturing protoplasts and subsequent plant regeneration from
10 cultured protoplasts. The second method requires (a) that the intact plant tissues, such as cotyledons, can be transformed by *Agrobacterium* and (b) that the transformed cells or tissues can be induced to regenerate into whole plants.

Most dicot species can be transformed by *Agrobacterium* as all species which are a natural plant host for *Agrobacterium* are transformable *in vitro*.
15 Monocotyledonous plants, and in particular, cereals, are not natural hosts to *Agrobacterium*. Attempts to transform them using *Agrobacterium* have been unsuccessful until recently (Hooykas-Van Slogteren *et al.*, *Nature* 311: 763-764 (1984)). However, there is growing evidence now that certain monocots can be transformed by *Agrobacterium*. Using novel experimental approaches cereal species

such as rye (de la Pena *et al.*, *Nature* 325: 274-276 (1987)), maize (Rhodes *et al.*, *Science* 240: 204-207 (1988)), and rice (Shimamoto *et al.*, *Nature* 338: 274-276 (1989)) may now be transformed.

Identification of transformed cells or plants is generally accomplished by
5 including a selectable marker in the transforming vector, or by obtaining evidence of successful bacterial infection.

Regeneration of Transformed Plants

Plant cells which have been transformed can also be regenerated using known techniques. For example, plant regeneration from cultured protoplasts is described in
10 Evans *et al.*, *Handbook of Plant Cell Cultures*, Vol. I: (MacMillan Publishing Co. New York, 1983); and I. R. Vasil (ed.), *Cell Culture and Somatic Cell Genetics of Plants*, (Acad. Press, Orlando, Vol. I, 1984, and Vol. II, 1986). It is known that practically all plants can be regenerated from cultured cells or tissues, including but not limited to, all major species of sugarcane, sugar beet, cotton, fruit trees, and
15 legumes.

Means for regeneration vary from species to species of plants, but generally a suspension of transformed protoplasts or a petri plate containing transformed explants is first provided. Callus tissue is formed and shoots may be induced from callus and subsequently rooted. Alternatively, somatic embryo formation can be induced in the
20 callus tissue. These somatic embryos germinate as natural embryos to form plants. The culture media will generally contain various amino acids and plant hormones,

such as auxin and cytokinins. It is also advantageous to add glutamic acid and proline to the medium, especially for such species as corn and alfalfa. Efficient regeneration will depend on the medium, on the genotype, and on the history of the culture. If these three variables are controlled, then regeneration is usually reproducible and

5 repeatable.

Silencing of Transgenes

It has been observed that many exogenously supplied expression units that are introduced into a plant are not expressed the level that might be expected based on the promoter and control sequences employed (M. Matzke, *et al.*, (1995)). The

10 inactivation of transgenes is known as silencing. Using the plants of the present invention, it has been observed that many introduced expression units become inactivated as a result of DNA methylation. Accordingly, the methods used to alter the amount of DNA methylation present in a plant, and plants generated through such methods, can be further used to control the level of expression of an introduced

15 expression unit.

The following examples are provided to illustrate, but not limit, the present invention. All references herein referred to are hereby incorporated by reference.

EXAMPLE 1

Plants With Reduced Rates of Maturation

1. Construction of antisense expression vectors

A 4.3 kb *MET1* cDNA spanning the positions indicated in Figure 1 was
5 inserted in the reverse orientation with respect to the CaMV 35S promoter in the
pMON530 T-DNA vector. *MET1* cDNA B5-2 was cloned into the *Eco RI* site of
T-DNA vector pMON530 (Monsanto) in the antisense orientation to the CaMV 35S
promoter through use of *Eco RI* sites present in cDNA linkers. The pMON530
T-DNA confers resistance to kanamycin (50 µg/mi) on transgenic plants.

10 2. Southern analysis of repetitive and single-copy DNA methylation patterns.

Total genomic DNA (3 µg/lane) from antisense lines, wild type, and the *ddm1*
mutant were digested with *Hpa II* (Fig. 2 left panels) or *Msp I* (Fig. 2 right panels),
subjected to
15 electrophoresis in 0.8% agarose, and transferred to Zeta-Probe membranes (Bio-Rad)
and hybridized as described to various probes (J. Chen, *et al.*, in *The Maize
Handbook*, M. Freeling, *et al.*, (Springer-Verlag, New York, 1994), pp. 525-527 and
S.L. Dellaporta, *et al.*, *ibid*, pp. 559-572). The filters were probed with a 180-bp
centromere repeat (J. M. Martinez-Zapater, *et al.*, (1985)) and 5S rDNA (B. R.
20 Campbell, *et al.*, (1992)) (Upper panels), or were probed with four single-copy gene

probes— PHOSPHORIBOSYLANTHRANILATE TRANSFERASE 1 (PAT1),
PROLIFERA (PRL), CONSTITUTIVELY PHOTOMORPHOGENIC 1 (COP1), and
ERECTA (ER) (Lower panels).

3. Southern and phenotypic analyses of Tr246 outcross progeny

5 Genomic DNAs (4 μ g/lane) from outcross progeny of strong antisense line
Tr246 to wild-type *Arabidopsis* (Columbia strain with no mutations) were digested
with *Hpa* II (Fig. 3 upper panels) or *Eco* RI (Fig. 3 lower panels). Southern analyses
were done as described above. Filters were probed with the 180-bp centromere repeat
(J. M. Martinez-Zapater, *et al.*, (1985)) (Fig. 3 upper panels) or the CaMV 35S
10 promoter fragment from pMON530 (Fig. 3 lower panels). Symbols indicate the
presence (+) or absence (-) of the antisense transgene. Phenotypic data for each
individual plant are shown below each lane. Plants were grown under continuous
light at 21°C.

4. Control of DNA Methylation Through Antisense Expression

15 To address the role of DNA methylation in plant development, an antisense
strategy was used to interfere with *MET1* a DNA methyltransferase (MTase) gene of
Arabidopsis, previously cloned by homology to the mouse gene (Fig. 1) (E. J.
Finnegan, *et al.*, (1993)). The *MET1* (cDNA B5-2) used in these experiments was
cloned independently; it lacked 436 base pairs (bp) downstream of the deduced

transaction start site and differed from the published sequence by a thymine-to-cytosine substitution at position +2,981. The *MET1* gene represents one member of a small gene family in *Arabidopsis* (E. J. Finnegan, *et al.*, (1993)) that maps to position 68.9 on chromosome 5, nonallelic to the *ddm1* locus (The *MET1* gene was mapped with an *Eco RI* polymorphism between *Arabidopsis* ecotypes WS and W100 with RI lines (Dupont) and is nonallelic to *ddm1* locus. T. Kakutani, *et al.*, (1995)). The *MET1* gene is expressed in seedling, vegetative, and floral tissues; in the inflorescence, expression is seen at highest levels in meristematic cells by *in situ* RNA hybridization.

10 To inhibit expression of the *MET1* gene, an antisense construct (Fig. 1), consisting of a 4.3-kb *MET1* cDNA in the antisense orientation under the control of a constitutive viral promoter (CaMV 35S), was introduced, into *Arabidopsis* strain Columbia. In one experiment, a total of nine primary transformants (TO generation) were recovered; progeny tests indicated that six lines that were further characterized
15 contained single-locus transferred DNA (T-DNA) insertions. All transformations and analyses used an *Arabidopsis* strain Columbia line homozygous for a mutation at the *g/l* locus as a marker (but otherwise wild type), except as noted by *Agrobacterium*-mediated transformation (N. Bechtold, *et al.*, *Acad. Sci. Paris* 315: 1,194 (1993)).

Six single-locus lines were identified by kanamycin segregation. Southern blot
20 analyses revealed that all lines contained independent insertions consisting of tandem repeats of the T-DNA. The plants used in this study represent the kanamycin-resistant

progeny of T1 outcrosses from single-locus lines Tr242, 243, 244, 245, and 248.

Before germination, all seeds were plated on MS media (Gibco) with (all transgenic lines) or without (controls) kanamycin (50 µg/ml, Sigma) and incubated for 3 days at 4°C in the dark. Growth conditions were 16 hours light, 8 hours dark at 21°C, except
5 as noted. Seedlings were transplanted into soil 8 days after germination.

Methylation patterns in repetitive DNA sequences were examined by Southern (DNA) hybridization (J. Chen, *et al.*, in *The Maize Handbook*, M. Freeling, *et al.*, (Springer-Verlag, New York, 1994), pp. 525-527 S.L. Dellaporta, *et al.*, *ibid*, pp. 559-572). Genomic DNAs were digested with the isochizomers *Hpa II* or *Msp I* (Fig.
10 2, upper panels) and probed with a centromeric repeat or a 5S ribosomal DNA (rDNA) sequence; both repeats are methylated in wild-type genomic DNA (A. Vongs, *et al.*, (1993)). *Hpa II* digestion is inhibited if either cytosine in the CCGG target site is methylated; *Msp I* can cleave C^{5m}CGC but not ^{5m}CCGG (M. Nelson, *et al.*, *Nucleic Acids Res.* 19: 2,045 (1991)).

15 With both probes, *Hpa II* digestion revealed a high extent of demethylation in three of six antisense lines (Tr244, 246, and 248; designated "weak") showed near wild-type levels of methylation. *Msp I* digestion was more complete in strong antisense lines contain substantial demethylation of these repeated sequences at C^{5m}CGG and ^{5m}CCGG sites.

20 Digestion of wild-type genomic DNA with *Eco RI* and *Bst NI*, isochizomers with differential sensitivity to cytosine methylation in the motif C^{5m}C(A/T)GG,

showed only minor differences in wild-type DNA when probed with the centromere repeat and 5S rDNA probes suggesting C(A/T)G methylation may not be prevalent.

DNA methylation was examined at four single-copy gene sequences (A. B. Rose, *et al.*, *Plant Physiol.* **100**: 582 (1992); P. S. Springer, *et al.*, *Science* **268**: 877 (1995); X. W. Deng, *et al.*, *Cell* **71**: 191 (1992); K. U. Torii, *et al.*, *Plant Cell* **8**: 735 (1995)) (Fig. 2, lower panels). Substantial demethylation of all four genes was seen only in the strong antisense lines; the *ddm1* mutant showed little or no demethylation relative to wild-type DNA, consistent with published reports (A. Vongs, *et al.*, (1993)). Total genomic levels of 5^mC were also measured in these lines by
10 high-performance liquid chromatography HPLC conditions were as described (C. W. Gehrke *et al.*, *J. Chromatogr.* **301**: 199 (1984), with the following exceptions: genomic DNAs were isolated from plants immediately after the onset of flowering, treated with 20 µg of ribonuclease A (Sigma) for 30 min. at 37°C, followed by passage through a Sepharose CL-5B (Pharmacia) spin column. Nucleosides were
15 resolved on a Varian Vista 5500 Liquid Chromatograph with a Rainin Dynamax 5 µm Spherical Microsorb C18 column (100 Å pore size, 4.5 mm inner diameter by 15 cm length) with a 20 min. isocratic gradient of 2.5% methanol, 50 mM KH₂PO₄ (pH 4.0), followed by a 10 min. linear gradient to 8.0% methanol, 50 mM KH₂PO₄ (pH 4.0). (Table I).

20 Cytosine methylation levels in wild-type (6.4% of total genomic cytosine) and the decrease in the *ddm1* mutant (75%) agree with previously published estimates

(A. Vongs, *et al.*, (1993)). Total genomic 5^mC content in strong lines Tr246 and Tr248 was reduced 71% relative to wild-type levels; the weak line Tr245 showed a 34% reduction. The decrease in 5^mC levels in the strong lines by a factor of 3.5 is comparable to reductions seen in the *ddm1* mutant and the MTase knockout mouse
5 (R. Li, *et al.*, *Cell* 59: 915 (1992)). Unlike the pattern of demethylation of the *ddm1* mutant, however, *MET1* antisense expression resulted in substantial demethylation of both repetitive DNA and single-copy gene sequences.

Cytosine methylation levels in *Arabidopsis* wild-type, *ddm1* mutant, and three antisense lines was determined by reversed-phase HPLC using the method of Gehrke
10 *et al.*, (1984), with the following exceptions: genomic DNAs were isolated from plants immediately after the onset of flowering, treated with 20 µg of ribonuclease A (Sigma) for 30 min. at 37°C, followed by passage through a Sepharose CL-5B (Pharmacia) spin column. Nucleosides were resolved on a Varian Vista 5500 Liquid Chromatograph with a Rainin Dynamax 5 µm Spherical Microsorb C18 column (100
15 Å pore size, 4.5 mm inner diameter by 15 cm length) with a 20 min. isocratic gradient of 2.5% methanol, 50 mM KH₂PO₄ (pH 4.0), followed by a 10 min. linear gradient to 8.0% methanol, 50 mM KH₂PO₄ (pH 4.0). All values presented in Table 1 represent the averages of two to four individual replicates and were calculated by integration of peak areas with Dynamax HPLC Method Manager v1.2 (Rainin). Percentages of total
20 5^mC content [$5^mC / (5^mC + C)$] are normalized for absorbance differences between cytosine and 5^mC.

Table 1 Line	Total 5 ^m C (%)	%WT levels	% Decrease
Wild type	6.38 ± 0.69	100	0
Tr245	4.24±0.59	66.5	33.5
5 Tr246	1.84±0.27	28.9	71.1
Tr248	1.83±0.16	28.7	71.3
<i>ddm1</i>	1.60±0.04	25.0	75.0

Quantitative aspects of vegetative and inflorescence traits are given for wild-type Columbia strain and six antisense lines in Table 2. Flowering time is measured in days after germination. Flowering time refers to the number of days elapsed from seed germination until emergence of an inflorescence bolt 0.5 to 1.0 cm in height; leaf number refers to the number of vegetative leaves initiated before emergence of the primary inflorescence; and secondary branches refers to the number of branches initiated on the primary inflorescence axis. Values are calculated from a minimum of four individual plants from each line. Pool sizes for each line were as follows: wild-type; TR242; TR245, n = 5; Tr243 and Tr244, n = 4; Tr245, n = 11; and Tr248, n = 15.

Table 2 Line	Flowering time	Leaf number	Secondary branches
Wild type	26.2±1.7	9.2±1.3	4.3±0.5
Tr242	24.5±2.1	10.0±1.4	4.5±1.1
5 Tr243	27.8±0.4	11.0±0.7	4.3±0.4
Tr244	47.7±4.0	34.0±5.7	20.3±4.9
Tr245	27.7±7.5	11.5±1.5	4.3±0.8
Tr246	45.9±3.6	32.5±3.2	20.7±1.4
Tr248	46.3±4.7	32.9±2.7	20.5±2.0

10 Normal patterns of development were perturbed in strong antisense lines.

Under long day conditions, six single-locus lines were identified by kanamycin segregation. Southern blot analyses revealed that all lines contained independent insertions consisting of tandem repeats of the T-DNA. The plants used in this study represent the kanamycin-resistant progeny of T1 outcrosses from single-locus lines

15 Tr242, 243, 244, 245, and 248. Before germination, all seeds were plated on MS media (Gibco) with (all transgenic lines) or without (controls) kanamycin (50 µg/ml, Sigma) and incubated for 3 days at 4°C in the dark. Growth conditions were 16 hours light, 8 hours dark at 21°C, except as noted. Seedlings were transplanted into soil 8 days after germination; Tr244, Tr246 and Tr248 plants initiated 30 to 35 vegetative

nodes with delayed abaxial trichome production and flowered after 45 to 48 days (Table 2); these phenotypes were fully penetrant in all T-DNA-containing progeny of the strong antisense lines.

After a transition from vegetative to reproductive development, wild-type plants initiated a primary inflorescence axis with two to five secondary inflorescence branches subtended by cauline leaves, followed by an abrupt transition to the production of solitary floral meristems (Table 2) (S. Shannon, *et al.*, *Plant Cell* 5: 539 (1993)). In strong antisense lines, the primary inflorescence shoot produced an average of 20 secondary branches (Table 2) before the production of flowers. The basal-most branches in strong antisense lines often assumed characteristics of vegetative rosettes, including enhanced spiral phyllotaxy of 10 to 20 vegetative-like leaves and shortened internodes, followed by the emergence of an inflorescence bolt recapitulating the primary inflorescence. Unlike wild-type plants, occasional secondary branches were produced apically to flowers in the transition zone. Early initiating flowers from strong antisense lines were normal in appearance and male-fertile, but were often female-sterile. Apical flowers initiated on secondary and tertiary inflorescences showed gross abnormalities, including a threefold increase in stamen number and sterile, incompletely fused carpels lacking stigmas. Despite the severe disruptions in floral morphology seen in these late-initiating flowers, no

obvious defects in pollen viability or paternal transgene transmission were observed on the basis of segregation ratios of transgenes in outcrosses. Kanamycin resistance segregated in a 1:1 ratio in all outcrosses from strong antisense lines.

One trivial explanation for the *MET1* antisense pleiotropy is that it represents an indirect effect on the transgene rather than a direct consequence of genomic demethylation; however, several lines of evidence support a direct role for DNA methylation in development. Data collected and analyzed in additional experiments revealed an identical pleiotropy in 12 independent transgenic lines. In analyses of outcross progeny from strong antisense lines, the severe phenotype co-segregated with the presence of the transgene, and a slightly attenuated pleiotropy was seen in progeny that had lost the transgene but retained a demethylated genome (Fig. 3). The latter finding is not unexpected, because the rate of genomic remethylation is slow (A. Vongs, *et al.*, (1993)). Phenotypic revertants seen among outcross progeny had reestablished near wild-type levels of genomic methylation (Fig. 3, lanes 5 and 11). In sum, it appears that demethylation is sufficient to maintain developmental pleiotropy in the absence of the transgene, whereas genomic remethylation is required to restore a wild-type phenotype.

The demethylation phenotype produced by antisense inhibition of *Arabidopsis* MTase differs markedly from the phenotype produced by the treatment of *Arabidopsis* seeds with the nonmethylatable cytosine analog 5-azacytidine (5-azaC), but bears some similarity to the *ddm1* mutant phenotype. 5-AzaC produces early flowering in

some late-flowering strains and mutants of *Arabidopsis* (J. E. Burn, *et al.*, *Proc. Nat'l Acad. Sci. U.S.A.* 90: 287 (1993)), but this effect has not been correlated with a quantitative decrease in cytosine methylation. More recent evidence suggests that the primary effect of 5-azaC treatment is due to toxicity resulting from covalent trapping
5 of DNA MTase (R. Jüttermann, *et al.*, *Proc. Nat'l Acad. Sci. USA* 91: 11,797 (1994)) or 5-azaC incorporation into RNA. The overall level of demethylation in the *ddm1* mutant is equivalent to that seen in the strong *MET1* antisense lines, yet the phenotype exhibited by the *MET1* antisense lines is much more severe and pleiotropic. This discrepancy may be due to substantial demethylation observed at single-copy gene
10 sequences in *MET1* antisense lines not seen in the *ddm1* mutant.

On the basis of these studies, DNA methylation is shown to be an essential component in the process of phase transitions and meristem determinacy. Methylation may serve as a primary signal to restrict meristem determinacy, or it may represent a secondary process required to maintain an epigenetic state once
15 established. Phase transitions involve an interplay of both cell autonomous and diffusible signals (R. S. Poethig, *Science* 250: 923 (1990)). Evidence that methylation may represent an autonomous component in this process comes from the observation that phase transitions are often irregular in strong antisense plan — the location of branches apical to flowers in the inflorescence transition zone may represent an
20 autonomous switching of individual cells in the meristem resulting in developmental mosaicism. Methylation effects have also been shown to be progressive in the plant

meristem (R. Martienssen, *et al.*, *Genes Dev.* 4: 331(1990)), and it is intriguing to speculate that a methylation gradient might be established during meristem growth to serve as an autonomous signal that directs meristem determinacy. In this light, the strong antisense phenotype could be explained by delayed establishment of this

5 hypothetical gradient. Such a model predicts that meristem potential becomes progressively more epigenetically restrictive and that repression cascades will be an underlying theme in plant determinacy, a process implicit in the pathway controlling inflorescence and floral development in *Arabidopsis* (M. D. Wilkinson, *et al.*, *Plant Cell* 7: 1,485 (1995)).

10

EXAMPLE 2

Plants with increased rates of development

To generate plants with increased amount of methylated DNA, *Arabidopsis* plants were altered to contain an expression unit that expresses cytosine methyl transferase under the control of the CaMV 355 promoter or another plant promoter.

15 The cDNA encoding the MET1 protein from *Arabidopsis* described above was used to generate expression units in which the *MET1* encoding sequence was placed under the regulatory control of the CaMV 355 promoter or the flower promoter AP3.

The expression units were introduced into a plant using *Agrobacterium* as described above. Transformed tissues were identified and plants were regenerated

20 from the transformed tissue.

An analysis of the rate of development was performed and it was found that plants having an increased amount of methylated DNA produced flowers at 10 to 12 days following germination, compared to 26 days for non-altered plants.

EXAMPLE 3

5 General methods for decreasing the rate of development in a plant

The Description of the Invention above provides a detailed outline for the various methods herein disclosed for altering the amount of DNA methylation present in a plant. The Examples provide the results of applications of the methods herein described. The following is intended to provide a non-limiting summary of the
10 typical steps employed when altering a plant using the present methods.

1. Select Plant

The present methods can be applied to any plant that can be altered using molecular techniques. As discussed above, there are numerous commercially exploited plants whose value would be increased by either shortening the time
15 required for maturation into seed producing plants or where it is desirable to increase biomass or seed yield. The first step in applying the present methods is to select as plant. Although it is preferable to select a plant that has had methods for expression, transformation and regeneration developed, most plants, though applied effort, can be transformed and regenerated using methods known in the art.

As part of choosing a plant, one needs to choose whether to increase or decrease the rate of maturation. As provided above, to increase the rate of maturation, the plant is altered so as to contain an increased amount of methylated DNA while to decrease the rate of maturation, the plant is altered so as to contained a decreased
5 amount of methylated DNA.

2. Chose the alteration method/target that is to be employed

The second step may be to decide which method to employ to alter the plant.

As provided above, the amount of DNA methylation can be decreased by inactivating or decreasing the activity of the methylase gene while the amount of DNA
10 methylation can be increased by activating or increasing the methylase activity within the cell.

selected based on the level of DNA methylation, the expression of the transgene, physiologic effects and equivalent methods thereto.

It should be understood that the foregoing discussion and examples merely present a detailed description of certain preferred embodiments. It therefore should be
5 apparent to those of ordinary skill in the art that based on the above description of the invention and the specific Examples provided, a skilled artisan can readily adapt the present invention for use with any plant that can be altered by using molecular techniques.. All articles and texts that are identified above are incorporated by reference in their entirety.

CLAIMS

1. A method to alter the rate of maturation in a plant comprising the step of genetically altering said plant using molecular techniques so that said plant has an altered amount of methylated DNA when compared to a non-altered plant sufficient to alter the rate of maturation of said plant.

2. The method of claim 1 wherein said alteration leads to a reduction in the amount of methylated DNA and said plant matures at a slower rate than said non-altered plant.

3. The method of claim 2, wherein said alteration results in about a 10% to about a 25% decrease, or greater, in the amount of methylated DNA present in said plant.

4. The method of claim 2, wherein said alteration results in about a 25% to about a 50% decrease, or greater, in the amount of methylated DNA present in said plant.

5. The method of claim 2, wherein said alteration results in about a 50% or about a 70% decrease, or greater, in the amount of methylated DNA present in said plant.

6. The method of claim 2, wherein said alteration leads to a reduction in the amount of methylated cytosine bases.

7. The method of claim 2 wherein said plant has had the activity of one or more DNA methyl transferase encoding genes altered using said molecular techniques.

8. The method of claim 7 wherein said DNA methyl transferase encoding gene is the *MET1* gene of *Arabidopsis*, or a homologue thereof.

9. The method of claim 2 wherein said plant has been altered using said molecular techniques so as to express an antisense nucleic acid molecule that is complementary to an mRNA that encodes a DNA methyl transferase.

10. The method of claim 9 wherein said mRNA is encoded by the *MET1* gene of *Arabidopsis*, or a homologue thereof.

11. The method of claim 1 wherein said alteration leads to an increase in the amount of methylated DNA and said plant matures at a faster rate than said non-altered plant.

12. The method of claim 11, wherein said alteration results in about a 10% to about a 25% increase, or greater, in the amount of methylated DNA present in said plant.

13. The method of claim 11, wherein said alteration results in about a 25% to about a 50% increase, or greater, in the amount of methylated DNA present in said plant.

14. The method of claim 11, wherein said alteration results in about a 50% to about a 70% increase, or greater, in the amount of methylated DNA present in said plant.

15. The method of claim 11, wherein said alteration leads to an increase in the amount of methylated cytosine bases.

16. The method of claim 15 wherein said plant has had one or more DNA methyl transferase encoding genes introduced into said plant using said molecular techniques.

17. The method of claim 16 wherein said DNA methyl transferase encoding gene is the *MET1* gene of *Arabidopsis*, or a homologue thereof.

18. A plant that has been altered according to the method of claim 1.
19. A plant that has been altered according to the method of claim 2.
20. A plant that has been altered according to the method of claim 11.
21. A method to increase the biomass production of a given plant comprising the step of genetically altering said plant using molecular techniques so that said plant has a reduced amount of methylated DNA when compared to a non-altered plant sufficient for said plant to mature at a slower rate than said non-altered plant.
22. The method of claim 21 wherein said plant has had the activity of one or more DNA methyl transferase encoding genes altered using said molecular techniques.
23. The method of claim 22 wherein said DNA methyl transferase encoding gene is the *MET1* gene of *Arabidopsis*, or an equivalent gene thereof.

24. The method of claim 21 wherein said plant has been altered using said molecular techniques so as to express an antisense nucleic acid molecule that is complementary to an mRNA that encodes a DNA methyl transferase.

25. The method of claim 21 wherein said mRNA is encoded by the *MET1* gene of *Arabidopsis*, or a homologue thereof.

26. A method to decrease the generation times in a plant breeding program comprising the step of using one or more parent strains that have been genetically altered using molecular techniques so that said parent strains have an increased amount of methylated DNA when compared to a non-altered plant sufficient to cause said parent strain to mature at a faster rate than said non-altered plant.

27. The method of claim 26 wherein said plant has had one or more DNA methyl transferase encoding genes introduced into said plant using said molecular techniques.

28. The method of claim 27 wherein said DNA methyl transferase encoding gene is the *MET1* gene of *Arabidopsis*, or a homologue thereof.

29. A method to select a plant that has an altered rate of development comprising the step of selecting a plant based on the criteria of having an altered amount of methylated DNA or an altered DNA methylase activity when compared to a non-altered plant sufficient to alter the rate of maturation of said plant.

30. The method of claim 29 wherein said alteration leads to a reduction in the amount of methylated DNA and said plant matures at a slower rate than said non-altered plant.

31. The method of claim 30, wherein said alteration results in about a 10% to about a 25% decrease, or greater, in the amount of methylated DNA present in said plant.

32. The method of claim 30, wherein said alteration results in about a 25% to about a 50% decrease, or greater, in the amount of methylated DNA present in said plant.

33. The method of claim 30, wherein said alteration results in about a 50% or about a 70% decrease, or greater, in the amount of methylated DNA present in said plant.

34. The method of claim 29 wherein said alteration leads to an increase in the amount of methylated DNA and said plant matures at a faster rate than said non-altered plant.

35. The method of claim 34, wherein said alteration results in about a 10% to about a 25% increase, or greater, in the amount of methylated DNA present in said plant.

36. The method of claim 34, wherein said alteration results in about a 25% to about a 50% increase, or greater, in the amount of methylated DNA present in said plant.

37. The method of claim 34, wherein said alteration results in about a 50% to about a 70% increase, or greater, in the amount of methylated DNA present in said plant.

38. A plant that has selected according to the method of claim 29.




FIG. 1




FIG. 2A

FIG. 2B





FIG. 2C **FIG. 2D** **FIG. 2E** **FIG. 2F** **FIG. 2G**

**FIG. 3**

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/13358

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C12N15/82 C12N9/10 C12N15/54 C12Q1/68 C12N5/10 A01H5/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C12N A01H		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	FINNEGAN, E.J.: "THE ROLE OF DNA METHYLATION IN PLANT DEVELOPMENT" EPIGENETIC MECHANISMS OF GENE REGULATION, vol. 32, 1996, COLD SPRING HARBOUR LABORATORY PRESS, pages 127-140, XP002049579 see the whole document ---	1-10, 18, 19, 21-25, 29-33, 38
X	KAKUTANI, T., ET AL.: "CHARACTERIZATION OF AN ARABIDOPSIS THALIANA DNA HYPOMETHYLATION MUTANT" NUCLEIC ACID RESEARCH, vol. 23, no. 1, 1995, pages 130-137, XP002049118 cited in the application PAGE 130; PAGE 132, RIGHT COLUMN, TABLE 1 --- -/--	1-6, 18, 19, 21, 29, 38
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.		
<input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
9 December 1997		23/12/1997
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Holtorf, S

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/13358

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	VONGS, A., ET AL. : "ARABIDOPSIS THALIANA DNA METHYLATION MUTANTS" SCIENCE, vol. 260, 25 July 1993, pages 1926-1928, XP002049119 cited in the application see the whole document ---	1,18,29, 38
A	FINNEGAN, E.J., ET AL. : "ISOLATION AND IDENTIFICATION BY SEQUENCE HOMOLOGY OF A PUTATIVE CYTOSINE METHYLTRANSFERASE FROM ARABIDOPSIS THALIANA" NUCLEIC ACID RESEARCH, vol. 21, no. 10, 1993, pages 2383-2388, XP002049120 cited in the application see the whole document ---	1-38
A	WO 96 14414 A (JOHN INNES CENTRE ;COUPLAND GEORGE MICHAEL (GB); PUTTERILL JOANNA) 17 May 1996 see the whole document ---	1-38
A	DE 44 44 460 A (INST GENBIOLOGISCHE FORSCHUNG) 30 May 1996 see the whole document ---	1-38
P,X	RONEMUS, M.J., ET AL. : "DEMETHYLATION-INDUCED DEVELOPMENTAL PLEIOTROPY IN ARABIDOPSIS" SCIENCE, vol. 273, 2 August 1996, pages 654-657, XP002049121 see figure 3; tables 1,2 ---	1-10,18, 19, 21-25, 29-33,38
P,X	FINNEGAN, E.J., ET AL. : "REDUCED DNA METHYLATION IN ARABIDOPSIS THALIANA RESULTS IN ABNORMAL PLANT DEVELOPMENT" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE USA. vol. 93, August 1996, pages 8449-8454, XP002049122 PAGE 8451, RIGHT COLUMN; PAGE 8452, RIGHT COLUMN; ABSTRACT ---	1,18,19, 29,38
T	RICHARDS, E.J., ET AL. : "DNA METHYLATION AND PLANT DEVELOPMENT" TRENDS IN GENETICS, vol. 13, no. 8, August 1997, pages 319-323, XP002049123 see the whole document -----	1-38

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/13358

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9614414 A	17-05-96	AU 3809795 A EP 0789765 A	31-05-96 20-08-97
DE 4444460 A	30-05-96	AU 4177096 A WO 9617069 A EP 0797673 A	19-06-96 06-06-96 01-10-97